arXiv:math/0002102v1 [math.AG] 13 Feb 2000

A $W(E_6)$-equivariant projective embedding of the moduli space of cubic surfaces

Masaaki YOSHIDA *

22 December 1999

Abstract

An explicit projective embedding of the moduli space of marked cubic surfaces is given. This embedding is equivariant under the Weyl group of type E_6. The image is defined by a system of linear and cubic equations. To express the embedding in a most symmetric way, the target would be 79-dimensional, however the image lies in a 9-dimensional linear subspace.

1 The moduli space of cubic surfaces

1.1 The space M and the action of the group $\underline{G}$

We first fix some notation and recall a few known facts on the moduli space of marked cubic surfaces. The moduli space of marked cubic surfaces, which we denote by M, is studied for example in [5] and [8]. Since any nonsingular cubic surface can be obtained by blowing up the projective plane $\mathbf{P}^2$ at six points, it can be represented by a 3×6-matrix of which columns give homogeneous coordinates of the six points. In order to get a smooth cubic surface from six points, we assume that no three points are collinear and the six points are not on a conic. On the set of 3×6 matrices, we have a cannical action of GL_3 on the left and the group $\mathbf{C}^\times$ acts naturally on homogeneous coordinates. By killing such ambiguity of coordinates, we get the following expression

$$x = \begin{pmatrix} 1 & 0 & 0 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 & x_1 & x_2 \\ 0 & 0 & 1 & 1 & x_3 & x_4 \end{pmatrix};$$

in this paper we use local coordinates (x_1, x_2, x_3, x_4) on M. The six points represented by the matrix above produces a non-singular cubic surface if and only if the following quantitiy does not vanish.

$$\begin{aligned} D(x) \quad := \quad & x_1x_2x_3x_4(x_1-1)(x_2-1)(x_3-1)(x_4-1) \\ & \times(x_1-x_2)(x_1-x_3)(x_2-x_4)(x_3-x_4)D_1D_2Q, \end{aligned}$$

---

*Department of Mathematics, Kyushu University, Fukuoka 812 Japan

where

$$\begin{aligned} D_1 &:= x_1x_4 - x_2x_3, \\ D_2 &:= x_1x_4 - x_4 + x_2 - x_2x_3 + x_3 - x_1, \\ Q &:= -x_2x_3x_1 - x_2x_3x_4 + x_2x_3 + x_1x_4x_2 + x_1x_4x_3 - x_1x_4, \end{aligned}$$

Thus we can identify the moduli space M with the affine open set $\{x = (x_1, \ldots, x_4) \mid D(x) \neq 0\}$.

Let us define as in [8] six bi-rational transformations $s_1, \ldots, s_6$ in $x = (x_1, \ldots, x_4)$:

$$\begin{aligned} s_1 : (x_1, x_2, x_3, x_4) &\to \left(\frac{1}{x_1}, \frac{1}{x_2}, \frac{x_3}{x_1}, \frac{x_4}{x_2}\right), \\ s_2 : (x_1, x_2, x_3, x_4) &\to (x_3, x_4, x_1, x_2), \\ s_3 : (x_1, x_2, x_3, x_4) &\to \left(\frac{x_1 - x_3}{1 - x_3}, \frac{x_2 - x_4}{1 - x_4}, \frac{x_3}{x_3 - 1}, \frac{x_4}{x_4 - 1}\right), \\ s_4 : (x_1, x_2, x_3, x_4) &\to \left(\frac{1}{x_1}, \frac{x_2}{x_1}, \frac{1}{x_3}, \frac{x_4}{x_3}\right), \\ s_5 : (x_1, x_2, x_3, x_4) &\to (x_2, x_1, x_4, x_3), \\ s_6 : (x_1, x_2, x_3, x_4) &\to \left(\frac{1}{x_1}, \frac{1}{x_2}, \frac{1}{x_3}, \frac{1}{x_4}\right). \end{aligned}$$

If M is regarded as the configuration space of six points in $\mathbf{P}^2$, the transformation s_1, for example, corresponds to the interchange of the two points represented by the first two column vectors of the matrix x. Each s_i turns out to be a bi-regular involution on M, and they form a group $\underline{G}$ isomorphic to the Weyl group of type E_6; relation of the generators are given by the Coxeter graph



1.2 Root system Δ of type E_6

We review the root system Δ of type E_6, following [6]. Consider an 8-dimensional Euclidean space $\tilde{E}$ with a standard basis $\varepsilon_1, \cdots, \varepsilon_8$. Let $\langle \cdot, \cdot \rangle$ be the inner product on $\tilde{E}$ defined by $\langle \varepsilon_j, \varepsilon_k \rangle = \delta_{jk}$ and let E be the linear subspace of $\tilde{E}$ spanned by the six vectors

$$\varepsilon_1, \cdots, \varepsilon_5,\ \tilde{\varepsilon} = \varepsilon_6 - \varepsilon_7 - \varepsilon_8.$$

We introduce the 36 vectors:

$$r = -\frac{1}{2}(\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4 + \varepsilon_5 + \tilde{\varepsilon}),$$

$$r_{1j} = -\varepsilon_{j-1} + r_0, \quad 2 \leq j \leq 6$$

$$r_{jk} = \varepsilon_{j-1} - \varepsilon_{k-1}, \quad 2 \leq j < k \leq 6$$

$$r_{1jk} = -\varepsilon_{j-1} - \varepsilon_{k-1}, \quad 2 \le j < k \le 6$$
$$r_{ijk} = -\varepsilon_{i-1} - \varepsilon_{j-1} - \varepsilon_{k-1} + r_0, \quad 2 \le i < j < k \le 6$$

where $r_{ij} = r_{ji}, r_{ijk} = r_{jik} = r_{ikj}$,

$$r_0 = \frac{1}{2}(\varepsilon_1 + \varepsilon_2 + \varepsilon_3 + \varepsilon_4 + \varepsilon_5 - \tilde{\varepsilon}).$$

Note that

$$r \perp r_{ij}, \quad r_{ij} \perp r_{kl}, \quad r_{ij} \perp r_{ijk}, \quad r_{ij} \perp r_{klm}, \quad r_{ijk} \perp r_{ilm}.$$

The set

$$\Delta = \{\pm r,\ \pm r_{ij},\ \pm r_{ijk}\}$$

forms a root system of type E_6. For example,

$$r_{12}, \quad r_{123}, \quad r_{23}, \quad r_{34}, \quad r_{45}, \quad r_{56}$$

can serve as a system of positive simple roots; its extended Dynkin diagram is given as



The set $\{r,\ r_{jk},\ r_{ijk}\}$ is the totality of positive roots of Δ.

Let s_r, s_{ij} and s_{ijk} be the reflections on E with respect to r, r_{ij} and r_{ijk}. These reflections act on Δ as

$$\begin{array}{ll} s_r: & r_{ij} \leftrightarrow r_{ij}, \quad r_{ijk} \leftrightarrow r_{lmn}, \quad \{i,j,k,l,m,n\} = \{1,\ldots,6\}, \\ s_{ij}: & \text{permutation of the indices } i \text{ and } j, \end{array}$$

$$\begin{array}{lllllllllll} s_{123}: & r_{12} & \leftrightarrow & r_{12}, & r_{14} & \leftrightarrow & r_{234}, & r_{56} & \leftrightarrow & r_{56}, \\ & r_{145} & \leftrightarrow & r_{145}, & r & \leftrightarrow & r_{456}, & & & \end{array}$$

modulo signs. Let us define two reflection group

$$G_1 = \langle s_{12}, s_{23}, s_{34}, s_{45}, s_{56} \rangle \cong S_6, \quad G = \langle G_1, s_{123} \rangle \cong W(E_6),$$

where S_6 is the symmetric group on six numerals $\{1, \ldots, 6\}$, $W(E_6)$ the Weyl group of type E_6, and $\langle a, b, \ldots \rangle$ denotes the group generated by $a, b, \ldots$ Note that

$$G_1 \subset \langle G_1, s_r \rangle = G_1 \times \langle s_r \rangle \subset G,$$

and G acts transitively on Δ.

1.3 Naruki's cross-ratio variety

A smooth compactification of M known as Naruki's cross-ratio variety $\mathcal{C}$ ([5], [8]), embedded in $(\mathbf{P}^1)^{45}$, is the union of M and the 76 divisors. The 36 of them correspond to the positive roots of Δ (they are said to be of the first kind), the other 40 divisors (said to be of the second kind, and are isomorphic to $(\mathbf{P}^1)^3$) can be represented as follows: Take three subsets Δ_1, Δ_2, Δ_3 of Δ satisfying the following conditions:

- Each of Δ_1, Δ_2, Δ_3 is a root system of type A_2.
- Δ_1, Δ_2, Δ_3 are mutually orthogonal.
- The vectors in $\Delta_1 \cup \Delta_2 \cup \Delta_3$ span E.

Note that each one of such three root systems determines the other two.

Such a triple $\{\Delta_1, \Delta_2, \Delta_3\}$ dertermines a divisor. According to the naming of the roots, we must use two different expressions: The first one is of the form

$$\{\pm r_{12}, \pm r_{23}, \pm r_{13}\}, \quad \{\pm r_{45}, \pm r_{56}, \pm r_{46}\}, \quad \{\pm r, \pm r_{123}, \pm r_{456}\},$$

(the corresponding divisor is denoted by $Z_{123,456}$ in [8],) and the second one is of the form

$$\{\pm r_{12}, \pm r_{234}, \pm r_{134}\}, \quad \{\pm r_{34}, \pm r_{356}, \pm r_{456}\}, \quad \{\pm r_{56}, \pm r_{125}, \pm r_{126}\},$$

(the corresponding divisor is denoted by $Z_{12,34,56}$ in [8]). Note that

$$Z_{123,456} = Z_{456,123}, \quad Z_{12,34,56} = Z_{56,12,34} \neq Z_{12,56,34}.$$

Thus, permuting the indices under G_1, we have 40 $(= 10 + 30)$ such divisors. The group G acts transitively on these 40 divisors.

Remark: These divisors (of the second kind) are disjoint to each other, and can be blown-down to points. In fact they correspond bijectively to the cusps of the modular group studied in [1] (see also [3] and [7]).

2 Embedding $\varphi : M \to \mathbf{P}^{80-1}$

2.1 Coordinates on $\mathbf{P}^{80-1}$ and the action of G

Let $\mathcal{A}$ be the set of 40 labels $(123, 456)$ and $(12, 34, 56)$ with the following identification

$$\begin{aligned} (123, 456) &= (213, 456) = (132, 456) = (456, 123), \\ (12, 34, 56) &= (21, 34, 56) = (56, 12, 34). \end{aligned}$$

Since G acts on the set Δ of roots, it acts also on the set of 40 divisors above, and so that it also acts on the set $\mathcal{A}$ of 40 labels.

We introduce 80 homogeneous coordinates

$$y_\alpha, \quad y_{-\alpha}, \qquad \alpha \in \mathcal{A}$$

on $\mathbf{P}^{80-1}$. I define an action of G on $\mathbf{P}^{80-1}$ by the following action of the generators $s_{12}, \ldots, s_{56}$ and s_{123} on the coordinates. Let s be one of the generators and $\alpha \in \mathcal{A}$; we assign

$$\begin{array}{rcll} s(y_\alpha) & = & y_\alpha, \quad s(y_{-\alpha}) = y_{-\alpha} & \quad \text{if } s\alpha = \alpha, \\ s(y_\alpha) & = & y_{-\beta}, \quad s(y_{-\alpha}) = y_\beta & \quad \text{if } s\alpha = \beta \neq \alpha. \end{array}$$

2.2 Definition of φ

In this section we define a map $M \to \mathbf{P}^{80-1}$. For a 3×6 matrix $x = (x_{ij})$, we consider 80 polynomials of degree 18 as follows:

$$\begin{array}{rcl} y_{(123,456)}(x) & = & D_{123}(x)D_{456}(x)Q(x), \\ y_{(12,34,56)}(x) & = & D_{134}(x)D_{234}(x)D_{356}(x)D_{456}(x)D_{512}(x)D_{612}(x), \end{array}$$

and $y_{-\alpha}(x) = -y_\alpha$ ($\alpha \in \mathcal{A}$), where $Q(x)$ is the determinant of the 6×6-matrix with columns

$$(x_{1j}x_{2j}, x_{2j}x_{3j}, x_{3j}x_{1j}, x_{1j}^2, x_{2j}^2, x_{3j}^2) \quad j = 1, \ldots, 6.$$

Since we have

$$y_\alpha(gxh) = (\det g)^6 y_\alpha(x)(\det h)^3,$$

the correswpondence above defines a map $\varphi : M \to \mathbf{P}^{80-1}$. For later use, we present 40 polynomials $y_\alpha(x)$ in terms of the coordinates (x_1, x_2, x_3, x_4) introduced in §1; the remaining 40 polynomoals are given by $y_{-\alpha}(x) = -y_\alpha(x)$. In the following table, $y_\alpha(x)$ is denoted simply by α, and we number them as $y_1, \ldots, y_{40}$:

$$\begin{array}{rcl}
y_1 = (156, 234) & := & D_1 Q : \quad y_2 = (123, 456) := D_2 Q : \\
y_3 = (124, 356) & := & (x_2 - x_1)Q : \quad y_4 = (145, 236) := (x_3 - x_1)Q : \\
y_5 = (146, 235) & := & (x_4 - x_2)Q : \quad y_6 = (134, 256) := (x_4 - x_3)Q : \\
y_7 = (135, 246) & := & x_1(x_4 - 1)Q : \quad y_8 = (136, 245) := x_2(x_3 - 1)Q : \\
y_9 = (125, 346) & := & x_3(x_2 - 1)Q : \quad y_{10} = (126, 345) := x_4(x_1 - 1)Q : \\
y_{11} = (12, 56, 34) & := & D_1(x_1 - 1)(x_2 - 1)(x_4 - x_3) : \\
y_{12} = (16, 23, 45) & := & D_1(x_1 - 1)(x_3 - 1)(x_4 - x_2) : \\
y_{13} = (15, 23, 46) & := & D_1(x_2 - 1)(x_4 - 1)(x_3 - x_1) : \\
y_{14} = (13, 56, 24) & := & D_1(x_3 - 1)(x_4 - 1)(x_2 - x_1) : \\
y_{15} = (15, 24, 36) & := & -D_1 x_1(x_2 - 1)(x_3 - 1) : \\
y_{16} = (16, 24, 35) & := & -D_1 x_2(x_1 - 1)(x_4 - 1) : \\
y_{17} = (15, 34, 26) & := & -D_1 x_3(x_1 - 1)(x_4 - 1) : \\
y_{18} = (16, 34, 25) & := & -D_1 x_4(x_2 - 1)(x_3 - 1) : \\
y_{19} = (12, 36, 45) & := & D_2 x_2 x_3(x_1 - 1) :
\end{array}$$

$$
\begin{aligned}
y_{20} = (12,35,46) &\;:=\; D_2x_1x_4(x_2-1) : \\
y_{21} = (13,26,45) &\;:=\; D_2x_1x_4(x_3-1) : \\
y_{22} = (13,25,46) &\;:=\; D_2x_2x_3(x_4-1) : \\
y_{23} = (13,24,56) &\;:=\; D_2x_1x_2(x_4-x_3) : \\
y_{24} = (15,46,23) &\;:=\; D_2x_1x_3(x_4-x_2) : \\
y_{25} = (16,45,23) &\;:=\; D_2x_2x_4(x_3-x_1) : \\
y_{26} = (12,34,56) &\;:=\; D_2x_3x_4(x_2-x_1) : \\
y_{27} = (13,46,25) &\;:=\; x_1(x_2-1)(x_3-1)(x_4-x_2)(x_4-x_3) : \\
y_{28} = (13,45,26) &\;:=\; x_2(x_1-1)(x_4-1)(x_3-x_1)(x_4-x_3) : \\
y_{29} = (12,46,35) &\;:=\; x_3(x_1-1)(x_4-1)(x_2-x_1)(x_4-x_2) : \\
y_{30} = (12,45,36) &\;:=\; x_4(x_2-1)(x_3-1)(x_2-x_1)(x_3-x_1) : \\
y_{31} = (14,35,26) &\;:=\; -x_1(x_1-1)(x_4-x_2)(x_4-x_3) : \\
y_{32} = (14,36,25) &\;:=\; -x_2(x_2-1)(x_3-x_1)(x_4-x_3) : \\
y_{33} = (14,25,36) &\;:=\; -x_3(x_3-1)(x_2-x_1)(x_4-x_2) : \\
y_{34} = (14,26,35) &\;:=\; -x_4(x_4-1)(x_2-x_1)(x_3-x_1) : \\
y_{35} = (16,25,34) &\;:=\; -x_2x_3(x_1-1)(x_4-x_2)(x_4-x_3) : \\
y_{36} = (15,26,34) &\;:=\; -x_1x_4(x_2-1)(x_3-x_1)(x_4-x_3) : \\
y_{37} = (16,35,24) &\;:=\; -x_1x_4(x_3-1)(x_2-x_1)(x_4-x_2) : \\
y_{38} = (15,36,24) &\;:=\; -x_2x_3(x_4-1)(x_2-x_1)(x_3-x_1) : \\
y_{39} = (14,56,23) &\;:=\; D_1D_2 : \\
y_{40} = (14,23,56) &\;:=\; (x_2-x_1)(x_3-x_1)(x_4-x_2)(x_4-x_3) :
\end{aligned}
$$

2.3 G-Equivariance of φ

Recall that the group $\underline{G}$ acts on M, and that G acts on $\mathbf{P}^{80-1}$. Let us identify the groups $\underline{G}$ and G by

$$\iota : s_{12} \mapsto s_1, \ldots, s_{56} \mapsto s_5,\ s_{123} \mapsto s_6.$$

Then we have

Theorem 1 *The map $\varphi : M \to \mathbf{P}^{80-1}$ is G-equivariant:*

$$g(\varphi(x)) = \varphi(\iota(g)x), \quad g \in G,\ x \in M,$$

that is,

$$(gy_\alpha)(x) = c_g y_\alpha(\iota(g)x), \quad g \in G,\ \alpha \in \pm\mathcal{A},\ x \in M,$$

where c_g is a rational function in (x_1, x_2, x_3, x_4).

Convention: Once this theorem is established, we ignore the redundant ones $y_{-\alpha}(x) = -y_\alpha(x)$ and regard φ as the map

$$M \ni x \longmapsto : y_\alpha(x) :\in \mathbf{P}^{40-1}.$$

The group G still acts on $\mathbf{P}^{40-1}$ by the transformations given in§2.3.

In order to prove the theorem, we have only to check the identity for a set of gnerators of G. Under s_1, the fourty polynomials are transformed as follows:

$$\begin{array}{lllll}
y_1 \to -c_1 y_6, & y_2 \to c_1 y_2, & y_3 \to c_1 y_3, & y_4 \to -c_1 y_8, & y_5 \to -c_1 y_7, \\
y_6 \to -c_1 y_1, & y_7 \to -c_1 y_5, & y_8 \to -c_1 y_4, & y_9 \to c_1 y_9, & y_{10} \to c_1 y_{10}, \\
y_{11} \to c_1 y_{11}, & y_{12} \to -c_1 y_{28}, & y_{13} \to -c_1 y_{27}, & y_{14} \to -c_1 y_{40}, & y_{15} \to -c_1 y_{32}, \\
y_{16} \to -c_1 y_{31}, & y_{17} \to -c_1 y_{35}, & y_{18} \to -c_1 y_{36}, & y_{19} \to c_1 y_{19}, & y_{20} \to c_1 y_{20}, \\
y_{21} \to -c_1 y_{25}, & y_{22} \to -c_1 y_{24}, & y_{23} \to -c_1 y_{39}, & y_{24} \to -c_1 y_{22}, & y_{25} \to -c_1 y_{21}, \\
y_{26} \to c_1 y_{26}, & y_{27} \to -c_1 y_{13}, & y_{28} \to -c_1 y_{12}, & y_{29} \to c_1 y_{29}, & y_{30} \to c_1 y_{30}, \\
y_{31} \to -c_1 y_{16}, & y_{32} \to -c_1 y_{15}, & y_{33} \to -c_1 y_{38}, & y_{34} \to -c_1 y_{37}, & y_{35} \to -c_1 y_{17}, \\
y_{36} \to -c_1 y_{18}, & y_{37} \to -c_1 y_{34}, & y_{38} \to -c_1 y_{33}, & y_{39} \to -c_1 y_{23}, & y_{40} \to -c_1 y_{14},
\end{array}$$

where $c_1 = (x_1 x_2)^{-3}$, under s_2,

$$\begin{array}{lllll}
y_1 \to y_1, & y_2 \to y_2, & y_3 \to -y_6, & y_4 \to y_4, & y_5 \to y_5, \\
y_6 \to -y_3, & y_7 \to -y_9, & y_8 \to -y_{10}, & y_9 \to -y_7, & y_{10} \to -y_8, \\
y_{11} \to -y_{14}, & y_{12} \to y_{12}, & y_{13} \to y_{13}, & y_{14} \to -y_{11}, & y_{15} \to -y_{17}, \\
y_{16} \to -y_{18}, & y_{17} \to -y_{15}, & y_{18} \to -y_{16}, & y_{19} \to -y_{21}, & y_{20} \to -y_{22}, \\
y_{21} \to -y_{19}, & y_{22} \to -y_{20}, & y_{23} \to -y_{26}, & y_{24} \to y_{24}, & y_{25} \to y_{25}, \\
y_{26} \to -y_{23}, & y_{27} \to -y_{29}, & y_{28} \to -y_{30}, & y_{29} \to -y_{27}, & y_{30} \to -y_{28}, \\
y_{31} \to -y_{33}, & y_{32} \to -y_{34}, & y_{33} \to -y_{31}, & y_{34} \to -y_{32}, & y_{35} \to -y_{37}, \\
y_{36} \to -y_{38}, & y_{37} \to -y_{35}, & y_{38} \to -y_{36}, & y_{39} \to y_{39}, & y_{40} \to y_{40},
\end{array}$$

under s_3,

$$\begin{array}{lllll}
y_1 \to c_3 y_1, & y_2 \to -c_3 y_3, & y_3 \to -c_3 y_2, & y_4 \to -c_3 y_7, & y_5 \to -c_3 y_8, \\
y_6 \to c_3 y_6, & y_7 \to -c_3 y_4, & y_8 \to -c_3 y_5, & y_9 \to c_3 y_9, & y_{10} \to c_3 y_{10}, \\
y_{11} \to c_3 y_{11}, & y_{12} \to -c_3 y_{16}, & y_{13} \to -c_3 y_{15}, & y_{14} \to -c_3 y_{39}, & y_{15} \to -c_3 y_{13}, \\
y_{16} \to -c_3 y_{12}, & y_{17} \to c_3 y_{17}, & y_{18} \to c_3 y_{18}, & y_{19} \to -c_3 y_{29}, & y_{20} \to -c_3 y_{30}, \\
y_{21} \to -c_3 y_{34}, & y_{22} \to -c_3 y_{33}, & y_{23} \to -c_3 y_{40}, & y_{24} \to -c_3 y_{38}, & y_{25} \to -c_3 y_{37}, \\
y_{26} \to c_3 y_{26}, & y_{27} \to -c_3 y_{32}, & y_{28} \to -c_3 y_{31}, & y_{29} \to -c_3 y_{19}, & y_{30} \to -c_3 y_{20}, \\
y_{31} \to -c_3 y_{28}, & y_{32} \to -c_3 y_{27}, & y_{33} \to -c_3 y_{22}, & y_{34} \to -c_3 y_{21}, & y_{35} \to c_3 y_{35}, \\
y_{36} \to c_3 y_{36}, & y_{37} \to -c_3 y_{25}, & y_{38} \to -c_3 y_{24}, & y_{39} \to -c_3 y_{14}, & y_{40} \to -c_3 y_{23},
\end{array}$$

where $c_3 = (1-x_3)^{-3}(1-x_4)^{-3}$, under s_4,

$$\begin{array}{lllll}
y_1 \to -c_4 y_5, & y_2 \to c_4 y_2, & y_3 \to -c_4 y_9, & y_4 \to c_4 y_4, & y_5 \to -c_4 y_1, \\
y_6 \to -c_4 y_7, & y_7 \to -c_4 y_6, & y_8 \to c_4 y_8, & y_9 \to -c_4 y_3, & y_{10} \to c_4 y_{10}, \\
y_{11} \to -c_4 y_{29}, & y_{12} \to c_4 y_{12}, & y_{13} \to -c_4 y_{40}, & y_{14} \to -c_4 y_{27}, & y_{15} \to -c_4 y_{33}, \\
y_{16} \to -c_4 y_{35}, & y_{17} \to -c_4 y_{31}, & y_{18} \to -c_4 y_{37}, & y_{19} \to c_4 y_{19}, & y_{20} \to -c_4 y_{26}, \\
y_{21} \to c_4 y_{21}, & y_{22} \to -c_4 y_{23}, & y_{23} \to -c_4 y_{22}, & y_{24} \to -c_4 y_{39}, & y_{25} \to c_4 y_{25}, \\
y_{26} \to -c_4 y_{20}, & y_{27} \to -c_4 y_{14}, & y_{28} \to c_4 y_{28}, & y_{29} \to -c_4 y_{11}, & y_{30} \to c_4 y_{30}, \\
y_{31} \to -c_4 y_{17}, & y_{32} \to -c_4 y_{38}, & y_{33} \to -c_4 y_{15}, & y_{34} \to -c_4 y_{36}, & y_{35} \to -c_4 y_{16}, \\
y_{36} \to -c_4 y_{34}, & y_{37} \to -c_4 y_{18}, & y_{38} \to -c_4 y_{32}, & y_{39} \to -c_4 y_{24}, & y_{40} \to -c_4 y_{13},
\end{array}$$

where $c_4 = (x_1x_3)^{-3}$, under s_5,

$$\begin{array}{lllll}
y_1 \to y_1, & y_2 \to y_2, & y_3 \to y_3, & y_4 \to -y_5, & y_5 \to -y_4, \\
y_6 \to y_6, & y_7 \to -y_8, & y_8 \to -y_7, & y_9 \to -y_{10}, & y_{10} \to -y_9, \\
y_{11} \to y_{11}, & y_{12} \to -y_{13}, & y_{13} \to -y_{12}, & y_{14} \to y_{14}, & y_{15} \to -y_{16}, \\
y_{16} \to -y_{15}, & y_{17} \to -y_{18}, & y_{18} \to -y_{17}, & y_{19} \to -y_{20}, & y_{20} \to -y_{19}, \\
y_{21} \to -y_{22}, & y_{22} \to -y_{21}, & y_{23} \to y_{23}, & y_{24} \to -y_{25}, & y_{25} \to -y_{24}, \\
y_{26} \to y_{26}, & y_{27} \to -y_{28}, & y_{28} \to -y_{27}, & y_{29} \to -y_{30}, & y_{30} \to -y_{29}, \\
y_{31} \to -y_{32}, & y_{32} \to -y_{31}, & y_{33} \to -y_{34}, & y_{34} \to -y_{33}, & y_{35} \to -y_{36}, \\
y_{36} \to -y_{35}, & y_{37} \to -y_{38}, & y_{38} \to -y_{37}, & y_{39} \to y_{39}, & y_{40} \to y_{40},
\end{array}$$

and under s_6,

$$\begin{array}{lllll}
y_1 \to -c_6y_{39}, & y_2 \to c_6y_2, & y_3 \to -c_6y_{26}, & y_4 \to -c_6y_{25}, & y_5 \to -c_6y_{24}, \\
y_6 \to -c_6y_{23}, & y_7 \to -c_6y_{22}, & y_8 \to -c_6y_{21}, & y_9 \to -c_6y_{20}, & y_{10} \to -c_6y_{19}, \\
y_{11} \to c_6y_{11}, & y_{12} \to c_6y_{12}, & y_{13} \to c_6y_{13}, & y_{14} \to c_6y_{14}, & y_{15} \to -c_6y_{18}, \\
y_{16} \to -c_6y_{17}, & y_{17} \to -c_6y_{16}, & y_{18} \to -c_6y_{15}, & y_{19} \to -c_6y_{10}, & y_{20} \to -c_6y_9, \\
y_{21} \to -c_6y_8, & y_{22} \to -c_6y_7, & y_{23} \to -c_6y_6, & y_{24} \to -c_6y_5, & y_{25} \to -c_6y_4, \\
y_{26} \to -c_6y_3, & y_{27} \to c_6y_{27}, & y_{28} \to c_6y_{28}, & y_{29} \to c_6y_{29}, & y_{30} \to c_6y_{30}, \\
y_{31} \to -c_6y_{35}, & y_{32} \to -c_6y_{36}, & y_{33} \to -c_6y_{37}, & y_{34} \to -c_6y_{38}, & y_{35} \to -c_6y_{31}, \\
y_{36} \to -c_6y_{32}, & y_{37} \to -c_6y_{33}, & y_{38} \to -c_6y_{34}, & y_{39} \to -c_6y_1, & y_{40} \to c_6y_{40},
\end{array}$$

where $c_6 = (x_1x_2x_3x_4)^{-2}$. Maybe it is interesting to see what happens under the operation of the involution s_r (classically called the association) which sends (x_1, x_2, x_3, x_4) to

$$\left(\frac{(x_4-1)D_1}{(x_4-x_2)(x_4-x_3)}, \frac{(x_3-1)D_1}{(x_3-x_1)(x_4-x_3)}, \frac{(x_2-1)D_1}{(x_4-x_2)(x_2-x_1)}, \frac{(x_1-1)D_1}{(x_3-x_1)/(x_2-x_1)}\right):$$

$$\begin{array}{lllll}
y_1 \to c_ry_1, & y_2 \to c_ry_2, & y_3 \to c_ry_3, & y_4 \to c_ry_4, & y_5 \to c_ry_5, \\
y_6 \to c_ry_6, & y_7 \to c_ry_7, & y_8 \to c_ry_8, & y_9 \to c_ry_9, & y_{10} \to c_ry_{10}, \\
y_{11} \to -c_ry_{26}, & y_{12} \to -c_ry_{25}, & y_{13} \to -c_ry_{24}, & y_{14} \to -c_ry_{23}, & y_{15} \to -c_ry_{38}, \\
y_{16} \to -c_ry_{37}, & y_{17} \to -c_ry_{36}, & y_{18} \to -c_ry_{35}, & y_{19} \to -c_ry_{30}, & y_{20} \to -c_ry_{29}, \\
y_{21} \to -c_ry_{28}, & y_{22} \to -c_ry_{27}, & y_{23} \to -c_ry_{14}, & y_{24} \to -c_ry_{13}, & y_{25} \to -c_ry_{12}, \\
y_{26} \to -c_ry_{11}, & y_{27} \to -c_ry_{22}, & y_{28} \to -c_ry_{21}, & y_{29} \to -c_ry_{20}, & y_{30} \to -c_ry_{19}, \\
y_{31} \to -c_ry_{34}, & y_{32} \to -c_ry_{33}, & y_{33} \to -c_ry_{32}, & y_{34} \to -c_ry_{31}, & y_{35} \to -c_ry_{18}, \\
y_{36} \to -c_ry_{17}, & y_{37} \to -c_ry_{16}, & y_{38} \to -c_ry_{15}, & y_{39} \to -c_ry_{40}, & y_{40} \to -c_ry_{39},
\end{array}$$

where

$$c_r = \left(\frac{D_1D_2}{(-x_4+x_2)(x_4-x_3)(x_1-x_3)(x_1-x_2)}\right)^3.$$

2.4 φ embeds M

It is known in [10] that the map

$$M \ni x \longrightarrow y_1(x) : y_3(x) : y_4(x) : y_5(x) : y_7(x) \in \mathbf{P}^4$$

is two-to-one, and induces an embedding of the quotient space $M/\langle s_r \rangle$. Thus the composite of φ and the projection

$$M \longrightarrow \mathbf{P}^{40-1} \longrightarrow \mathbf{P}^4$$

is a two-to-one map. This fact together with the equivariance of φ under the involution s_r shown just above implies

Theorem 2 *φ embeds M into $\mathbf{P}^{40-1}$.*

2.5 Prolongation of φ to degenerate arrangements

Let us consider degenerate arrangements of six points on the plane. Since arrangements with three collinear points can be transformed under G to those with six points on a conic, we assume, Without loss of generality, that our arrangements represented by $x = (x_1, \ldots, x_4)$ satisfies $Q = 0$, that is, the six points are on a conic. Since a (nonsingular) conic is isomorphic to a line, such arrangements form the configuration space

$$X(2,6) = GL(2) \backslash \{Mat(2,6) \mid \text{any } 2 \times 2 \text{ minor} \neq 0\} / (\mathbf{C}^\times)^6$$

of six points on the projective line: if we represent a point of $X(2,6)$ by a matrix of the form

$$z = \begin{pmatrix} 1 & 0 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & z_1 & z_2 & z_3 \end{pmatrix},$$

where

$$\prod_{i=1}^{3} z_i(z_i - 1) \prod_{1 \le i < j \le 3} (z_i - z_j) \neq 0,$$

then the degenerate arrangements in question can be parametrized by $z = (z_1, z_2, z_3)$ as

$$x_1 = (1 - z_1)/(1 - z_2), \quad x_2 = (1 - z_1)/(1 - z_3), \quad x_3 = z_1/z_2, \quad x_4 = z_1/z_3.$$

Among the two S_6-equivariant projective embedding of $X(2,6)$ presented in [9], let us recall the following one given by the fifteen polynomials

$$D_{ij}(z) D_{kl}(z) D_{mn}(z), \quad \{i, j, k, l, m\} = \{1, \ldots, 6\},$$

where $D_{ij}(z)$ is the (i,j)-minor of the 2×6-matrix z. Their actual forms are given by

$$(z_i - 1)(z_j - z_k), \quad z_j - z_k, \quad z_i(z_j - z_k), \quad z_i(z_j - 1).$$

It is known and easy to show that the image is projectively equivalent to the so-called Segre cubic defined by

$$t_0 + \cdots, t_5 = 0, \quad (t_0)^3 + \cdots + (t_5)^3 = 0.$$

On the other hand, let us prolong the domain of definition of the map φ on these degenerate arrangements by the same forty polynomials. Then the map φ in z-coodinates is given by $y_1 = \cdots y_{10} = 0$ and

$$\begin{array}{lll}
cy_{11} = -z_1(z_2 - z_3), & cy_{12} = -z_2 + z_1, & cy_{13} = z_1 - z_3, \\
cy_{14} = -(-1 + z_1)(z_2 - z_3), & cy_{15} = (-1 + z_1)z_3, & cy_{16} = (-1 + z_1)z_2, \\
cy_{17} = z_1(-1 + z_3), & cy_{18} = z_1(-1 + z_2), & cy_{19} = -(-z_2 + z_1)z_3, \\
cy_{20} = -(z_1 - z_3)z_2, & cy_{21} = -(-z_2 + z_1)(-1 + z_3), & cy_{22} = -(z_1 - z_3)(-1 + z_2), \\
cy_{23} = -(-1 + z_1)(z_2 - z_3), & cy_{24} = z_1 - z_3, & cy_{25} = -z_2 + z_1, \\
cy_{26} = -z_1(z_2 - z_3), & cy_{27} = -(z_1 - z_3)(-1 + z_2), & cy_{28} = -(-z_2 + z_1)(-1 + z_3), \\
cy_{29} = -(z_1 - z_3)z_2, & cy_{30} = -(-z_2 + z_1)z_3, & cy_{31} = (-1 + z_3)z_2, \\
cy_{32} = (-1 + z_2)z_3, & cy_{33} = (-1 + z_2)z_3, & cy_{34} = (-1 + z_3)z_2, \\
cy_{35} = z_1(-1 + z_2), & cy_{36} = z_1(-1 + z_3), & cy_{37} = (-1 + z_1)z_2, \\
cy_{38} = (-1 + z_1)z_3, & cy_{39} = z_2 - z_3, & cy_{40} = z_2 - z_3,
\end{array}$$

where

$$c = \frac{(z_1 - 1)(z_1 - z_3)(z_2 - z_3)(z_1 - z_2)z_1}{(1 - z_2)^2 z_3^2 (1 - z_3)^2 z_2^2}.$$

This shows that the prolonged φ gives exactly the embedding of the arranged arragements given by $\{x \mid Q(x) = 0\}$, isomorphic to $X(2, 6)$, given above.

Further put

$$x_1 = t\xi_1, \quad x_2 = t\xi_2, \quad x_3 = t\xi_3, \quad x_4 = 1 + t\xi_4,$$

and let t tends to zero in $\varphi(x)$. We can easily see that the limit is a point whchi is independ of $(\xi_1, \ldots, \xi_4)$.

These results together with the facts on Naruki's cross-ratio $\mathcal{C}$ variety reviewd in §1.3, we can readily show

Theorem 3 *The closure of the image of M under φ is isomorphic to the variety obtained from Naruki's cross-ratio $\mathcal{C}$ by blowing down the 40 exceptional divisors of the second kind to points.*

Remark: This variety is isomorphic to the Stake compactification of the modular variety obtained in [1]. Note that the 40 cusps correspond to the 40 points obtained by blowing down the divisors of the second kind.

3 Defining equations

We will find a set of generators of the ideal defining the closure $\overline{\varphi(M)}$ of the image of M in $\mathbf{P}^{40-1}$. Among the forty polynomials $y_1(x), \ldots, y_{40}(x)$, we can find some relations. The Prücker relations ([10]) of the 3×3-minors of a 3×6-matrix yield linear relations among $y_1(x), \ldots, y_{10}(x)$; for example,

$$(124, 356) - (145, 236) + (146, 235) - (134, 256) = y_3(x) - y_4(x) + y_5(x) - y_6(x) = 0.$$

On the other hand, it is easy to check the cubic relation

$$\begin{aligned}&(124,356)(15,23,46)(13,26,45)-(145,236)(13,56,24)(12,35,46)\\&=y_3(x)y_{13}(x)y_{21}(x)-y_4(x)y_{14}(x)y_{20}(x)=0.\end{aligned}$$

Let V be the subvariety of $\mathbf{P}^{40-1}$, coordinatized by $y_1,\dots,y_{40}$, defined by the G-orbits of the linear and the cubic equations:

$$y_3-y_4+y_5-y_6=0,\quad y_3y_{13}y_{21}-y_4y_{14}y_{20}=0.$$

Theorem 4 $\overline{\varphi(M)}=V$.

3.1 An outline of the proof

By operating the group G to the linear equation, we get many linear relations among $y_1,\dots,y_{40}$. These relations form a system of rank 30, that is, all y's can be expressed linearly in terms of ten chosen ones. For example, in terms of

$$y_1,\ y_3,\ y_4,\ y_5,\ y_7,\ y_{11},\ y_{12},\ y_{13},\ y_{15},\ y_{19},$$

the remaining thirty ones are expressed as

$$\begin{aligned}
y_2&=y_1-y_5+y_4,\quad y_6=-y_4+y_5+y_3,\quad y_8=-y_3-y_1+y_7,\\
y_9&=-y_1+y_7-y_4,\quad y_{10}=y_7-y_5-y_3,\quad y_{14}=y_{11}+y_{13}-y_{12},\\
y_{16}&=-y_1+y_{12}-y_{13}-y_{11}+y_{15},\quad y_{17}=y_{15}-y_1-y_{13},\\
y_{18}&=-y_{13}-y_{11}+y_{15},\quad y_{20}=y_{19}+y_3+y_{11},\quad y_{21}=y_4+y_{19}+y_{12},\\
y_{22}&=y_{19}+y_3+y_{11}+y_{13}+y_5,\quad y_{23}=y_1-y_{12}+y_{13}+y_3+y_{11},\\
y_{24}&=y_4+y_1+y_{13},\quad y_{25}=y_{12}+y_5-y_1,\\
y_{26}&=-y_4+y_5-y_1+y_3+y_{11},\quad y_{27}=y_{19}+y_4+y_1+y_3-y_7+y_{11}+y_{13},\\
y_{28}&=-y_7+y_{19}+y_{12}+y_5+y_3,\quad y_{29}=y_3-y_7+y_{11}+y_{19}+y_5,\\
y_{30}&=y_3+y_1-y_7+y_{19}+y_4,\quad y_{31}=-y_1-y_{13}+y_{19}+y_{15}+y_{12},\\
y_{32}&=y_{19}+y_3+y_{15},\quad y_{33}=y_{19}+y_4+y_{15},\\
y_{34}&=y_{19}-y_{13}+y_{15}+y_{12}+y_5-y_1+y_3,\\
y_{35}&=-y_5-y_1-y_3+y_7-y_{11}+y_{15}-y_{13},\\
y_{36}&=-y_1+y_7-y_4-y_{13}+y_{15},\\
y_{37}&=-y_3+y_7-y_{11}+y_{15}-y_1+y_{12}-y_{13},\\
y_{38}&=-y_1+y_7+y_{15},\quad y_{39}=y_1-y_{12}+y_{13},\\
y_{40}&=-y_5+y_4+y_1+y_{13}-y_{12}.
\end{aligned}$$

By operating the group G to the cubic equation, we get many such relations. Substituting the expressions of the y's obtained above, we get cubic relations in terms of the chosen ten y's; let us here rename the ten coordinates as:

$$\begin{aligned}&g_1=y_1,\ g_2=y_3,\ g_3=y_4,\ g_4=y_5,\ g_5=y_7,\\&g_6=y_{11},\ g_7=y_{12},\ g_8=y_{13},\ g_9=y_{15},\ g_0=y_{19}.\end{aligned}$$

Among these cubic equations we can find exactly thirty linearly independent ones. Therefore the variety V is isomorphic to a subvariety of $\mathbf{P}^9$, coodinatized by $g_1, \ldots, g_9, g_0$, defined by thirty cubic equations, say $cub_1, \ldots, cub_{30}$. Some of them which are relatively simple are shown below:

$$
\begin{aligned}
cub_1 \; := \;& g_2g_8g_0 + g_2g_8g_7 - g_3g_6g_0 - g_2g_3g_6 - g_3g_6^2 - g_8g_0g_3 - g_3g_6g_8 \\
& + g_3g_7g_0 + g_2g_3g_7 + g_6g_7g_3, \\
cub_2 \; := \;& g_0^2g_1 + g_3g_1g_0 + g_1^2g_0 + g_2g_0g_1 - g_5g_1g_0 + g_0g_1g_6 + g_8g_1g_0 + g_2g_3g_6 \\
& + g_2g_6g_1 + g_2g_8g_6 + g_3g_1g_6 + g_1^2g_6 + g_8g_1g_6 - g_5g_3g_6 - g_5g_1g_6 - g_5g_8g_6, \\
cub_3 \; := \;& -g_8g_0g_3 + g_8g_0g_4 + g_2g_8g_0 - g_3g_6g_0 - g_2g_3g_6 - g_3g_6^2 - g_3g_6g_8 - g_3g_6g_4, \\
cub_4 \; := \;& g_2g_0g_7 + g_2^2g_7 + g_2g_6g_7 + g_2g_8g_7 + g_2g_7g_4 - g_0g_4g_6 - g_8g_0g_4 + g_0g_4g_7, \\
cub_5 \; := \;& -g_5g_3g_2 - g_2g_5g_1 - g_5g_8g_2 + g_2g_3g_0 + g_2g_0g_1 + g_2g_8g_0 + g_2g_3g_7 \\
& + g_2g_1g_7 + g_2g_8g_7 + g_2g_3g_4 + g_2g_1g_4 + g_2g_8g_4 + g_2^2g_3 + g_2^2g_1 + g_2^2g_8 + g_5g_4g_0 \\
& - g_5g_3g_0 - g_5g_1g_0 - g_5g_8g_0 + g_5g_7g_0, \\
cub_6 \; := \;& g_5g_8g_1 - g_8g_1g_4 - g_2g_8g_1 - g_5g_8g_7 + g_4g_7g_8 + g_2g_8g_7 \\
& + g_5g_8^2 - g_8^2g_4 - g_2g_8^2 + g_5g_8g_2 - g_2g_8g_4 - g_2^2g_8 + g_5g_8g_6 - g_8g_4g_6 - g_2g_8g_6 \\
& - g_5g_7g_6 - g_5g_4g_6 + g_5g_1g_6, \\
cub_7 \; := \;& -g_9g_2g_3 + g_3g_9g_5 - g_3g_9g_6 - g_9g_3g_0 - g_9g_3g_4 + 2g_9g_2g_4 - g_4g_9g_5 \\
& + g_4g_9g_6 + g_9g_0g_4 + g_9g_4^2 + g_9g_2^2 - g_9g_5g_2 + g_9g_2g_6 + g_9g_0g_2 \\
& - g_5g_6g_0 - g_5g_3g_6 - g_5g_6g_9, \\
cub_8 \; := \;& -g_5g_3g_7 + g_5g_4g_7 + g_2g_5g_7 + g_5g_7g_6 - g_2g_3g_1 - g_2g_0g_1 - g_2g_1g_7 + g_5g_3g_1 \\
& + g_5g_1g_0 - g_3g_1g_6 - g_0g_1g_6 - g_1g_7g_6 - g_3g_1g_0 - g_0^2g_1 - g_1g_7g_0 \\
& - g_1g_4g_3 - g_1g_4g_0 - g_1g_4g_7, \\
cub_9 \; := \;& -g_8g_1g_3 - g_3g_1g_6 + g_3g_8g_7 + g_6g_7g_3 - g_9g_3g_7 - g_3g_8^2 - 2g_3g_6g_8 \\
& + g_9g_3g_8 - g_3g_2g_8 - g_2g_3g_6 + g_9g_2g_3 - g_3g_6^2 + g_3g_9g_6 - g_3g_1g_0 - g_2g_3g_1 \\
& - g_0^2g_1 - g_2g_0g_1 - g_9g_0g_1 - g_1g_7g_0 - g_2g_1g_7 - g_9g_1g_7, \\
cub_{10} \; := \;& g_5g_8g_0 - g_8g_0g_4 - g_2g_8g_0 + g_5g_8g_9 - g_9g_8g_4 - g_2g_9g_8 + g_3g_6g_8 \\
& + g_8g_0g_3 + g_2g_3g_6 - g_5g_3g_6 + g_3g_6^2 + g_3g_6g_0 + g_3g_6g_4 - g_9g_2g_3 + g_3g_9g_5 \\
& - g_3g_9g_6 - g_9g_3g_0 - g_9g_3g_4, \\
cub_{11} \; := \;& g_2g_5g_1 + g_5g_1^2 - g_5^2g_1 + g_5g_1g_0 + g_5g_3g_1 - g_2g_5g_7 - g_5g_1g_7 + g_5^2g_7 \\
& - g_5g_7g_0 - g_5g_3g_7 + g_5g_8g_2 + g_5g_8g_1 - g_5^2g_8 + g_5g_8g_0 + g_5g_3g_8 \\
& - g_3g_2g_8 - g_2g_8g_0 - g_2g_8g_7, \\
& \vdots \\
cub_{19} \; := \;& -g_5g_1g_0 - g_2g_5g_1 + g_0^2g_1 + 2g_2g_0g_1 + g_0g_1g_6 + g_1g_7g_0 + g_2g_1g_7 + g_1g_7g_6 \\
& + g_1g_4g_0 + g_2g_1g_4 + g_1g_4g_6 + g_2^2g_1 + g_2g_6g_1 - g_5g_7g_6 - g_5g_4g_6, \\
& \vdots
\end{aligned}
$$

We first study this system over the field $K := \mathbf{C}(g_1, \ldots, g5)$. Geometrically speaking, we project the variety V onto the 4-dimensional projective space coordinatized by $g_1 : \cdots : g_5$, and study the generic fibre of the projection

$$\pi : \mathbf{P}^9 \supset V \ni g_1 : \cdots : g_0 \longmapsto g_1 : \cdots : g_5 \in \mathbf{P}^4.$$

We shall prove that π is generically two-to-one. This implies π is two-to-one on

$$V^\circ := V - V \cap \cup_{j=1}^{40} \{y_j = 0\}.$$

Thus the argument in §2.4 shows that $\varphi : M \to V^\circ$ is an isomorphism.

We next study the intersection $V \cap \{y_1 = 0\}$, and prove that $V \cap \cup_{j=1}^{40} \{y_j = 0\}$ is the totality of the G-orbit of the closure of the image of $X(2,6)$ under the prolonged φ, stated in §2.5.

3.2 Computation over K

From $cub_3 = 0$ and $cub_7 = 0$, we can solve g_8 and g_9 as:

$$\begin{aligned} g_8 &= g_3g_6(g_2 + g_6 + g_0 + g_4)/(g_0g_2 - g_0g_3 + g_0g_4 - g_3g_6), \\ g_9 &= g_5g_6(g_0 + g_3)/(-g_2g_3 + g_5g_3 - g_3g_6 - g_0g_3 - g_4g_3 + 2g_2g_4 - g_5g_4 + g_4g_6 + g_0g_4 \\ &\quad + g_4^2 + g_2^2 - g_5g_2 + g_2g_6 + g_0g_2 - g_5g_6). \end{aligned}$$

Substituting these into cub_{19}, we can solve g_6:

$$g_6 = -\frac{g_1(-g_5g_0 - g_5g_2 + g_0^2 + 2g_0g_2 + g_0g_4 + g_7g_0 + g_2g_7 + g_2^2 + g_2g_4)}{g_0g_1 + g_1g_4 + g_1g_7 + g_2g_1 - g_5g_7 - g_5g_4}.$$

Substituting these expressions into $cub_1, \dots, cub_{30}$, we have $cub_3 = cub_7 = cub_{19} = 0$ of course and $cub_{10} = 0$; though most of the remaining ones are complicated, cub_8 is relatively simple:

$$cub_8 = -(g_1 - g_5)qq/(g_0g_1 + g_1g_4 + g_1g_7 + g_2g_1 - g_5g_7 - g_5g_4),$$

where

$$\begin{aligned} qq &= g_1g_4t^2 + (-g_2g_5 - g_5g_4 + g_1g_4 + g_5g_3)s^2 + 2g_1g_4st \\ &+ (-g_5g_3g_1 + g_2g_1g_5 + g_1g_4g_3 - g_4^2g_5 + g_1g_2g_4 + g_5g_3g_4 - g_2g_5g_4 + g_4^2g_1)s \\ &+ (-g_4g_5g_1 + g_4^2g_1 + g_1g_4g_3 + g_1g_2g_4)t + g_3g_2g_1g_4 - g_5g_3g_1g_4 + g_1g_4^2g_3 \end{aligned}$$

is a quadratic form in

$$t := g_0, \quad s := g_7.$$

By this quadratic equation, we reduce the degree with respect to t of the numerators of the cub's to 1. In this way we get 26 ($= 30 - 4$) equations of the form

$$cube_j : a_jt + b_j = 0, \quad j = 1, \dots, 30, \quad j \neq 3, 7, 10, 19,$$

where $a_j, b_j \in K[s]$. The polynomials

$$D_{jk} = a_jb_k - a_kb_j \in K[s], \quad j < k$$

have a unique common factor dd, which is quadratic in s; its actual form is given by

$$\begin{aligned}
dd \;:=\; & (g_1^2g_4^2+2g_2g_1^2g_4+g_3^2g_5^2+g_2^2g_3^2+g_2^2g_1^2+g_5^2g_4^2+2g_3g_2g_1g_4-2g_2g_1g_5g_4\\
& -2g_3g_5^2g_4-2g_1g_5g_4^2+2g_2^2g_3g_1-2g_3^2g_5g_2+2g_5g_3g_2g_4+2g_5g_3g_1g_4-2g_5g_3g_1g_2)s^2\\
+\; & (-g_2^2g_1^3-2g_1^3g_2g_4+2g_1^2g_2g_4^2+2g_1^2g_5g_4^2+g_1^2g_4^3-g_1^3g_4^2-2g_1g_5g_4^3\\
& -2g_3g_5^2g_4^2-2g_3g_2^2g_1^2+2g_3g_2g_1g_4^2+g_3^2g_2^2g_4+g_5^2g_4^3+2g_1^2g_2g_5g_4\\
& +2g_2^2g_3g_1g_4-4g_3g_5g_1g_2g_4+g_5^2g_3^2g_4+g_2^2g_1^2g_4-g_1g_5^2g_4^2-g_2^2g_3^2g_1\\
& -g_3^2g_5^2g_1+2g_5g_3g_2g_4^2+2g_5g_1^2g_2g_3+2g_3^2g_5g_1g_2-2g_1g_2g_5g_4^2\\
& +2g_3g_5^2g_1g_4+2g_3g_1g_5g_4^2-2g_3g_5g_1^2g_4-2g_3^2g_5g_2g_4-2g_1^2g_2g_3g_4)s\\
-\; & g_1^2g_3^2g_4^2+g_3g_1^2g_4^3-g_1^3g_3g_4^2-g_1^2g_2g_3^2g_4-g_1^3g_3g_2g_4-g_3^2g_1g_2^2g_4\\
& -g_3^2g_1g_2g_4^2+g_3g_1g_2g_4^3+g_2^2g_3g_1g_4^2-g_3g_5^2g_1^2g_4+g_1^3g_3g_5g_4\\
& +g_3^2g_5g_1g_4^2-g_3^2g_5^2g_1g_4+g_3g_5^2g_1g_4^2-g_3g_5g_1g_4^3-g_3g_1^2g_2^2g_4\\
& +g_3^2g_5g_1^2g_4-2g_3g_1g_2g_5g_4^2+2g_3g_1^2g_2g_5g_4+2g_3^2g_1g_2g_5g_4.
\end{aligned}$$

So $dd(s)=0$ is the compatibility condition of the system of equations $cube_j$: $a_j(s)t+b_j(s)=0$, among which $cube_{11}:a_{11}(s)t+b_{11}(s)=0$ has the minimal degree of coefficients; actually, degrees of a_{11} and b_{11} with respect to s are 3 and 4.

Now we get three equations $dd=0, qq=0$ and $cube_{11}=0$. Substituting the expression $t=-b_{11}/a_{11}$ into qq, we get a polynomial in s. We can prove that this polynomial has dd as a factor.

We have proved that for a given generic point $g_0:\cdots:g_5\in\mathbf{P}^4$, the inverse under π consists of two points. They are obtained as follows: solve the quadratic equation $dd=0$ with respect to $s(=g_7)$. For each root, $t(=g_0)$ is uniquely determined by the linear equation $cube_{11}$. And g_6,g_8,g_9 are uniquely determined by $cub3, cub7, cub19$ as stated above. Then all the relations are satisfied.

3.3 Intersection of V and $\{y_1=0\}$

We should better work on V living in $\mathbf{P}^{40-1}$. In terms of the forty coordinates $y_1,\dots,y_{40}$, the cubic equations are 2-term equations. Thus the vanishing of y_1 implies that of some other coordinates. Thanks to G-action, we can assume that $y_{10}=0$. The vanishing of these two coordinates forces the vanishing of other coordinates. Tedious case-by-case study shows that every component of $V\cap\{y_1=0\}$ is included in the G-orbit of

$$V\cap\{y_1=\cdots=y_9=y_0=0\},$$

which is the closure of the image of $\{x\mid Q(x)=0\}$ under the prolonged φ.

References

[1] D. Allcock, J. Carlson and D. Toledo, A complex hyperbolic structure for Moduli of cubic surfaces. C.R. Acad. Sci. 326(1998), 49–54.

[2] K. Matsumoto Theta constants associated with the triple covering of the complex projective line branching at 6 points, preprint, 1999, September.

[3] K. Matsumoto in preparation, 1999, September.

[4] K. Matsumoto, T. Sasaki and M. Yoshida: The monodromy of the period map of a 4-parameter family of K3 surfaces and the hypergeometric system of type (3,6). Intern. J. Math. **3** (1992), 1-164.

[5] I. Naruki, Cross ratio variety as a moduli space of cubic surfaces, Proc. london Math. Soc. 45(1982), 1–30

[6] B. Hunt, A remarkable quintic fourfold in $\mathbf{P}^5$ and its dual variety (private note 7.1.1992).

[7] T. Sasaki and M. Yoshida, The uniformizing differential equation of the complex hyperbolic structure on the moduli space of marked cubic surfaces, Proc. Japan Acad. **75**(1999), 129–133.

[8] J. Sekiguchi and M. Yoshida, $W(E_6)$-orbits of the configurations space of 6 lines on the real projective space, Kyushu J. Math. 51(1997), 1–58.

[9] M. Yoshida, The real loci of the configuration space of six points on the projective line and a Picard modular 3-fold, Kumamoto J. of Math. 11(1998), 43–67.

[10] M. Yoshida, *Hypergeometric Functions, My Love,* Vieweg Verlag, 1997.